



APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: **PROTEIN 68075 AND ITS USE FOR REGENERATING NERVE
CELL PROCESSES**

APPLICANT: **STUART A. LIPTON**

"EXPRESS MAIL" Mailing Label Number TB426550949US

Date of Deposit November 30, 1994

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as **"Express Mail Post Office To Addressee"** with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Meghan Madera
Meghan Madera



485-201

C.I.P.

A
08 346910

PATENT

ATTORNEY DOCKET NO: 00108/017004

PROTEIN 68075 AND ITS USE FOR
REGENERATING NERVE CELL PROCESSES

F

5 This application is a continuation under 37 CFR 1.78
of U.S.S.N. 08/167,109^{now abandoned} (hereby incorporated by reference),
filed December 14, 1993, which is a continuation of U.S.S.N.
07/932,926, filed August 20, 1992, now abandoned, which is a
continuation of U.S.S.N. 07/758,292, filed August 28, 1991,
10 now abandoned, which is a continuation of U.S.S.N.
07/619,653, filed November 27, 1990, now abandoned, which is
a continuation of U.S.S.N. 07/391,779, filed August 9, 1989,
now abandoned.

Background of the Invention

15 This invention relates to methods for enhancing
regeneration of mammalian nerve cells in vivo.

Summary of the Invention

In a first aspect, the invention features a purified
human protein, which we termed the 68075 protein, or a
20 purified protein comprising a portion of the 68075 protein
which is active in promoting regeneration of a process of a
central or peripheral neuron. By "68075 protein" is meant
the complete human protein having a portion encoded by the
cDNA clone deposited with the American Type Culture
25 Collection (ATCC) on August 9, 1989 and assigned the
Accession number 68075.

By "purified" is meant that the 68075 protein is
isolated from its natural environment, e.g., a human cell,
and one or more components with which it naturally occurs is
30 reduced in level in comparison to the natural environment.
It is preferred that 68075 be purified to the extent that it

forms at least 10% of the protein of a preparation, and even more preferably that it forms the majority of the protein in that preparation. When used for treatment of humans, it is preferred that the 68075 protein be in the form of a

5 homogeneous solution or lyophilized. That is, the 68075 protein is purified away from all compounds with which it naturally occurs. This term is meant to include purified naturally occurring 68075 polypeptide, but most preferably it refers to synthetic polypeptides forming the biologically
10 active portion of 68075 (i.e., that portion able to promote regeneration of a neuron process), and molecules formed by recombinant genetic engineering techniques. Thus, the term includes polypeptides expressed within a bacterial, mammalian, or other cell, from a strand of DNA which does
15 not naturally occur within that cell, or from DNA under the control of a regulatory sequence with which this DNA does not naturally occur.

The 68075 protein may be purified by a number of techniques. These include isolation of the naturally
20 occurring polypeptide by immunoaffinity procedures using antibodies against the polypeptide, or anti-idiotypes (antibodies against 68075 antibodies). They also include isolation and purification of cells which naturally express, or are modified to express, high levels of the 68075
25 polypeptide. Such cells may be formed by genetic engineering techniques to introduce the 68075 gene, and manipulating the engineered cell to cause expression of that gene within the cell. It is also possible to insert secretion sequences, well known to those skilled in the art,
30 upstream of the structural gene encoding 68075 to cause the polypeptide to be secreted into the medium surrounding the cells that then can be isolated from the medium.

The amino acid sequence of the part of the 68075 protein which is active in promoting regeneration of a process (e.g., an axon or a dendrite) of a central neuron may be determined by standard techniques, e.g., by
5 expressing short fragments of the DNA encoding 68075 and determining which of the resulting polypeptides are active. This sequence of amino acids can be synthesized by standard chemical procedures well known to those skilled in the art. Such recombinant and synthetic peptides are within this
10 invention.

Also included within the definition of polypeptide 68075 is 68075 which is caused to be produced in vivo within a human at a level at which it does not normally occur in that human cell. For example, agents which cause elevated
15 production of 68075 may be administered to the human cells in order to elevate the level of 68075 expressed from the naturally occurring gene. Such agents can be identified by standard procedure using the cloned 68075 encoding gene and investigating the effect of agents on expression of the gene
20 in vivo or in vitro. In addition, the naturally occurring gene may be modified by transfection of the cell with a virus in order to introduce a gene encoding a 68075, or an agent which modulates expression of the naturally occurring gene, to increase the level of expression of 68075 within
25 that cell.

In a second aspect, the invention features purified nucleic acid encoding human 68075. This nucleic acid is separated from the environment in which it naturally occurs and is generally located on a vector, for example, a
30 plasmid, cosmid, or phage, such that nucleic acid may be manipulated as desired. This nucleic acid includes not only nucleic acid which naturally occurs within a human cell, but modifications (made by standard procedures) of the nucleic

acid which cause production of a 68075 protein identical in amino acid sequence to a naturally occurring 68075, or having one or more amino acids substituted or deleted without significantly affecting the biological activity of

5 the resulting 68075 with regard to its promotion of the regeneration of a process of a central or peripheral neuron. Thus, the nucleotide bases within the nucleic acid may be substituted conservatively or non-conservatively using techniques well known to those skilled in the art.

10 Similarly, nucleic acid closely related to naturally occurring human nucleic acid is encompassed by this invention. Such nucleic acid may be identified by using standard probe screening techniques, where the probe is a portion of at least 15-20 contiguous bases of the nucleic

15 acid encoding human 68075. This probe nucleic acid is used under standard stringent hybridization conditions to identify nucleic acid homologous to that encoding the human 68075. Not all such homologous sequences will encode a 68075 protein, but those which do can be identified by

20 standard procedures.

In a third aspect the invention features a method for promoting regeneration of a process of a neuron of a human. The method features providing the purified or recombinant human 68075 described above, and applying that

25 protein to the neuron in an amount sufficient to promote regeneration of a process of a neuron of a human.

Some potential steps of providing and applying 68075 are briefly described above. These steps include provision of a homogenous preparation of 68075 protein directly to the

30 neuron to be treated, or application of any other equivalent protein as defined above. Neurons which may be treated in this invention include those of the central nervous system, as well as the peripheral nervous system. Such treatment

may include administering the substance in a nerve guide tube, well known to those skilled in the art, via an osmotic minipump, or by use of a slow release pellet, for example, an Elvax pellet. The protein may also be administered
5 intrathecally, stereotactically or intravitreally (for CNS), or systemically (for PNS).

Alternatively, the level of 68075 protein within a cell may be increased by regulation of the naturally occurring 68075 gene within that cell, or of a gene which is
10 introduced into the cell by a standard technique. For example, a viral promoter system (e.g., HSV-1 promoter) can be used to cause expression of 68075 in adult glial or other cells, e.g., nerve cells, which no longer, or do not naturally, express the 68075 protein, by insertion of that
15 promoter upstream of the natural 68075 encoding gene. In addition, a cell line of autologous cells (e.g., glia or other cell types) encoding 68075 can be transfected with such a viral promoter to cause expression of the protein and these cells transplanted into the CNS or PNS. Fragments of
20 cells or entire cells, e.g., young or passaged astrocytes that express 68075, can be introduced stereotactically into the nervous system.

Applicants have found that protein 68075 of mammals enhances the regeneration of nerve cell processes in vivo in
25 humans. These proteins are useful in treatment of diseases caused by nerve cell degeneration, for example, dementia or other injury, and also for treatment for peripheral nerve degeneration or severance. For example, after operations known to produce male impotence, administration of 68075
30 protein at the cut site of the neuron will enhance regeneration of the neuron.

Other features of the invention will be apparent from the following description of the preferred embodiments and from the claims.

Description of the Preferred Embodiments 68075 Protein

5 68075 protein has been defined above. There follows an example of isolation of clone 68075 from human cells. Those skilled in the art will recognize that this example is not limiting to the invention and other methods for isolation of purified 68075 will be readily apparent upon
10 reading of this disclosure. Similarly, the complete 68075 encoding gene can be readily isolated by standard techniques using the partial clones provided in this application. To this end, a deposit has been made of a clone including 500 bases encoding at least some of the 68075 polypeptide, and a
15 clone including 1340 bases encoding nearly all of the 68075 polypeptide. These deposits have been made in the American Type Culture Collection, ^{10801 University Boulevard, Manassas VA 20108-2209} ~~12301 Parklawn Drive, Rockville,~~
F F MD, under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms
20 for the purpose of Patent Procedure. Applicant's assignee, Children's Medical Center Corporation of Boston, represents that the ATCC is a depository affording permanence of the deposit and ready accessibility thereto by the public if a patent is granted. All restrictions on the availability to
25 the public of the material so deposited will be irrevocably removed upon the granting of a patent. The material will be available during the pendency of the patent application to one determined by the Commissioner to be entitled thereto under 37 CFR 1.14 and 35 USC 112. The deposited material
30 will be maintained with all the care necessary to keep it viable and uncontaminated for a period of at least five years after the most recent request for the furnishing of a sample of the deposited material, and in any case, for a

period of at least thirty years after the date of deposit or for the enforceable life of the patent, whichever period is longer. Applicant's assignee acknowledges its responsibility to replace the deposit should the depository
5 be unable to furnish a sample when requested due to the condition of the deposit. The first deposit is assigned the accession number 68075 and was deposited on August 9, 1989. The second deposit was made on November ³⁰, 1994 and is assigned the accession number, 75949 or 97525.

10 In the example to be described, anti-idiotypic antibodies to Thy-1 antibodies were used to screen a cDNA library of human DNA. The anti-idiotypic antibodies against Thy-1 antibodies were characterized by Pillemer et al. J. Experimental Medicine, 153:106, 1981, hereby
15 incorporated by reference. These were obtained and purified according to the methods described by Pillemer et al. These anti-idiotypic antibodies are not essential to the present invention. Applicant believes that the anti-idiotypic antibodies used in the present invention are available from
20 Pillemer et al, or commercially available from Sigma (St. Louis) or Hazelton Labs (PA).

The anti-idiotypic antibodies TEPC-15, HOPC-8, MOPC-603, and MOPC-167 of Pillemer et al. were purified from ascitic fluid over an anti-IgA affinity column. Media
25 surrounding the cells was centrifuge-filtered and passed through an Amicon concentrator. This was stored at -80° until use. The ascitic fluid was then passed over an anti-IgA affinity column, and the eluate dialyzed and concentrated against 0.5 M NaCl, 0.1 M NaHCO₃.

30 Example 1: cDNA encoding part of human 68075

A fetal human brain cDNA library was constructed in lambda gt11. To accomplish this, cDNA was synthesized from 10 mg of polyadenylated RNA by a modification of the method

of Gubler et al., Gene 25:263, 1983. A double-stranded cDNA was treated with *Eco*R1 methylase, ligated to *Eco*R1 linkers, cleaved with *Eco*R1, and passed over a Sepharose CL4-B column to remove excess linkers. The cDNA was ligated to lambda
5 gt11 DNA which had been pre-ligated, digested with 100-fold excess of *Eco*R1, and phosphatased using one unit of calf intestinal alkaline phosphatase (Boehringer Mannheim) per 70 mg of phage DNA. Approximately 500 ng of cDNA was ligated to 50 ng of lambda gt11 arms. The DNA was packaged
10 according to standard protocols and amplified in *E. coli* Y 1088 on tryptone plates (1% Bacto-tryptone, 0.5% NaCl, 1.2% agar, 2 mM MgCl₂) in tryptone top agar (0.8% agar). To screen the library, the phage was grown on Y1090 at a density of 100,000 per 150 mm plate. Standard protocol was
15 followed through incubation of the filters in the primary (anti-idiotypic) antibodies in Tris buffer with 1% goat serum. Filters were then washed twice in Tris buffer containing 0.1% Triton X-100, incubated in peroxidase-conjugated goat anti-mouse IgA second antibodies, washed in
20 50 mM Tris, pH 7.4, and visualized with diaminobenzidine and hydrogen peroxide. A positive plaque was then cloned and used in a Northern blot analysis for hybridization to mouse RNA obtained from cultures of astrocytes.

A preliminary screening of the library using the
25 anti-idiotypic antibodies TEPC-15 and HOPC-8 revealed a single positive clone consisting of approximately 500 base pairs of cDNA, originally designated TR1, now called 68075_{DNA}.

Rescreening of the brain library with the 500 base
30 clone identified two two kilobase (clones TR2A and TR2B (now accession number 75949 or 97525)) and three three kilobase (clones TR3A, TR3B, and TR3C) clones to which the 500 base clone hybridized. The mRNA detected on Northern

A G
Sub
1
I

Sub
I
blots with clone 68075_{DNA} is 6 kb in size. However, much of it may consist of nontranslated sequences. In Western blots, 30-60 kDa bands were detected, suggesting that 68075 requires only about two kilobases to encode the full length protein.

5 The above clone, 68075_{DNA}, is the cDNA included in ATCC Deposit No. 68075, and can be used to identify other candidate clones to insure that a full length 68075 clone is obtained. This clone may be manipulated by standard
10 procedures to determine the DNA sequence of the clone, and thence the amino acid sequence of 68075. Once these sequences are determined, the 68075 gene can be manipulated into any standard expression vector to cause expression of the 68075 protein from any number of standard expression
15 cells. The expressed 68075 protein is then isolated and purified by standard techniques. A 68075-encoding gene can also be transfected into mammalian cells, for example, by using recombinant clones in the DO vectors, such as DOL and/or DOJ, which contain the polyoma early region, or
20 pBR322. These retrovirus constructs are used to overexpress the protein in a mammalian cell line that does not normally express 68075. For example, NIH 3T3 cells can be used. In this method a sodium phosphate buffer at pH 6.95 is mixed with a calcium DNA solution, and the calcium phosphate DNA
25 precipitate formed allowed to precipitate gradually onto the cells to provide a high efficacy of stable transformation. Those cells transformed are expanded by standard procedure. An example of an eukaryotic expression library includes the pH3M vector, having an SV40 origin of replication, an M13
30 origin, a SupF marker, and a convenient cloning site. Cells expressing 68075 are selected by panning (e.g., by their adherence to dishes coated with anti-idiotypic antibody). DNA is isolated from adherent cells and used to transform

MC1061/p3 cells, amplified, and used to transfect COS cells for a second round of enrichment. Standard methods can be used to produce variants of the DNA clones, which may include variants of 68075 at the amino acid level.

5 Example 2: Affinity purification of 68075

68075 protein may be purified by affinity column purification. Either astrocyte cultures or whole brain preparations are solubilized in detergent (Nonidet P-40 or sodium deoxycholate), passed over the column, and then
10 equilibrated with buffer (1M Tris HCl, pH 7.4). Elution of 68075 is with acidic and basic buffers (e.g., 0.1 M glycine HCl, pH 2). The resulting fractions are assayed for protein, and those fractions containing protein assayed for 68075. 68075 is identified by standard SDS polyacrylamide
15 gel electrophoresis, or by immunoblot analysis. Alternatively, anti-idiotypic antibodies such as those discussed above, or monoclonal antibodies to 68075, may be used in its detection.

Methods

20 In order to use the above described 68075, any number of methods may be utilized as discussed above.

Example 3: Promotion of regeneration with 68075.

One method of this invention includes providing a supply of 68075 protein immobilized on a solid support.
25 This support is placed in proximity to a mammalian neural process in a living mammal to promote growth and regeneration of the process. The method of the invention is particularly applicable to neural processes including ganglion cell bodies, such as retinal ganglion cell bodies,
30 but can also be used to enhance regeneration of nerve cell processes in the peripheral as well as the central nervous system.

The term "solid support" includes both rigid materials such as glass, synthetic plastics, and the like on the surface of which a layer or coating of the monoclonal antibody is bonded, and also soft gels penetrable by the
5 regenerating nerve process such as collagen, fibrin, fibrinogen, blood clot, or laminin throughout which the monoclonal antibody is dispersed and bonded so that it is immobilized. A supply of 68075 immobilized in a soft gelatinous mass such as a proteinaceous gel, for example
10 collagen gel, enclosed within an open ended tube or hollow cylinder of glass or synthetic plastic which serves as a guide for the regenerating processes, the latter advancing through the center of the tube, can also be used. The synthetic plastic used for the tube or hollow cylinder can
15 be any nontoxic material to which antibodies bond, such as polyethylene, polypropylene, etc., but is preferably a bioresorbable material such as that described in Nyilas, U.S. Patent 4,481,353. The guide filled with proteinaceous gel on which 68075 is immobilized is implanted so as to
20 bridge the gap between the ends of a damaged or severed process to enhance regeneration and extension of the process through the center of the tube spaced from the synthetic plastic wall. The following example is intended to illustrate more fully the nature of the invention without
25 acting as a limitation upon its scope. It is understood that this example demonstrates the utility of 68075 in rats but that naturally or traumatically induced defects in neural processes can also be treated in this manner.

Male Sprague-Dawley rats can be utilized in these
30 studies. Four animals serve as an unoperated control and three animals receive an intracranial optic nerve transection alone. In eight animals, one optic nerve is transected intracranially and then a nerve guide tube

implanted to bridge the transected nerve. To achieve this, a small craniotomy is made in the frontal lobe and carefully aspirated to expose a 2 to 3-mm segment of one optic nerve intracranially. The pial plexus of capillaries together
5 with the anterior cerebral artery are gently retracted medially and the nerve transected. Such an intracranial nerve transection spares the blood supply to the retina.

After hemostasis is achieved, a 1.5-mm length (0.95 mm o.d, 0.75 mm i.d.) of the nerve guide is inserted. The
10 nerve guides used in these experiments are fabricated as polymers of synthetic poly D,L-lactates with 2% triethyl citrate as a plasticizer, as described in U.S. Patent 4,481,353.

The experimental animals are divided into three
15 groups. In one group (two rats), empty nerve guide tubes are implanted. In the second group (three rats), the nerve guides' lumen is filled with a collagen matrix (Vitrogen, Flow Laboratories, diluted in 0.1 M phosphate buffer to a final concentration of 2.48 mg/ml) containing a 0.5 mg/ml
20 fibrinogen (bovine, CalBiochem) immediately before implantation. The third group (three rats) receive nerve guides containing 68075, in addition to the collagen and fibrinogen.

In all animals, both the proximal and distal nerve
25 stumps are inserted into the guide. At the time the nerve guides are implanted, all experimental animals receive bilateral superior cervical ganglionectomies to remove the sympathetic innervation to the head. This procedure is necessary to rule out the ingrowth of peripheral sympathetic
30 fibers, a process which has been shown to occur after indirect and direct lesions to the CNS.

Four weeks postoperatively the animals are deeply anesthetized and perfused through the heart with 200 ml

heparinized saline, followed by 500 ml 1% paraformaldehyde plus 3% glutaraldehyde, followed by 200 ml 0.1 M phosphate buffer. All perfusion solutions are made up in 0.1 M phosphate buffer. The nerve guide is dissected out as one
5 piece, immersed in 1% osmium tetroxide, and processed for embedding in plastic (DER, Ted Pella, Inc.). One micrometer transverse sections are cut from the proximal, middle, and distal portions of the nerve guide and stained with toluidine blue. "Proximal" refers to sections across the
10 part of the nerve guide closest to the eye.

Data are collected with a computer-controlled light microscope at a final magnification of 1600X. Blood vessels are identified and their luminal profiles entered on-line from a digitizing tablet to a display terminal and then to a
15 VAX 11/780 computer (Digital Equipment Corp.) for further numerical and statistical analysis. For each cross-sectioned sample the tissue cable areas, the number of blood vessels, and the area of each blood vessel is obtained. This data allows determination of whether the 68075 is
20 active in promoting neural process growth.

This procedure can also be used to determine the use of various forms of 68075, or fragments thereof. For example, recombinant 68075 can be expressed by standard procedure and tested for its efficacy in this method. By
25 expression of parts of the 68075-encoding DNA that portion having the desired activity in this assay can readily be determined.

One other method for measuring the use of 68075 for neuronal regeneration is the retinal culture procedure of
30 Leifer et al, 1984, supra. Generally, rat and mouse retinal ganglion cells are identified with fluorescent dyes that are retrogradely transported to the ganglion cell bodies after injection into the superior colliculus and lateral

geniculate body, to which the ganglion cells project. Fluorescently-tagged Thy-1 antibodies are specific among retinal cells in vitro for the ganglion cells. Double labelling experiments show virtually complete overlap
5 between the two marking methods, ensuring the validity of either for identifying the ganglion cells. Ganglion cells are dissociated from the retinas of 4 day-old to adult rats or mice under anesthesia using papain (10 units/ml, Cooper Biomedical, Inc.) and mild mechanical trituration. The
10 dissociated retinal cells are plated in culture using minimal essential media and 5% rat serum (for rat cultures) or fetal calf serum (for mouse cultures). For rat cultures, Long-Evans rats are used, and, for mouse cultures, C57BL/6 mice.

15 To measure the regeneration of processes by the ganglion cells, retinal cells from a single dissociation are cultured with and without 68075 in multiple dishes (each receiving a 50 μ l drop of the same dilution of cells). On the second, third, and fourth days in culture, solitary
20 ganglion cells are measured for the diameter of the soma, and each process is measured using a computer graphics system.

Other Embodiments

Other embodiments are within the following claims.

What is claimed is: